




FIGURE 1**BEST AVAILABLE COPY**


FIGURE 2

006260" 8TH 24960

FIGURE 3



006260" 2744960

FIGURE 4

006260" 87424960



FIGURE 5

FIGURE 6

006260" 87424960

59/647418


FIGURE 7


FIGURE 8



FIGURE 9

FIGURE 10

006260" 8T424960

FIGURE 11

F _v 2-Casp-3						F _v 2-Casp3/S163					
AP1903 nM						AP1903 nM					
100	32	10	3.2	1	0	100	32	10	3.2	1	0

09/647418-092900


FIGURE 12



FIGURE 13

FIGURE 14

006260" 2744350



FIGURE 15

FIGURE 16



006260"BT44950




FIGURE 17

FIGURE 18



006260" 87424950

FIGURE 19



**B**

FIGURE 20

A



B




FIGURE 21

CID-Mediated Apoptosis In BPH-derived CR-2a SMC




FIGURE 22

CID-Mediated Apoptosis in BPH derived JD SMC




Fig 23 (Fig 24 on Reverse)






Fig 25 (Fig 26 on Reverse)



Final Size: 67%
or 7 cm

Fig 27



Final Size: 67%
or 6.9 cm


Generation of Ad-GFP-CMV-YM-E

Ad-GFP-CMV-E-ICE US 99/06799

pAdTrack-CMV ~ Fv1s1 - YM-E
 ~ E-Fv1-Fv1s1 - ICEst

① cut pAdTrack-CMV with EcoRV + Not I

1 μ g/2 μ l (pAdTrack-CMV) + 1 μ l Buffer H
 + 1 μ l EcoRV + 1 μ l Not I + H₂O 5 μ l.
 37°C, 2 hr.



pAdTrack-CMV
 9220 bp

② cut pSH1/s-Fv1s1-YM-E with Not I + EcoRI
 (got 2.2 kb Fv1s1-YM-E)

(E-Fv1-Fv1s1-ICEst)
 or

2 μ g/12 μ l (pSH1/s-Fv1s1-YM-E) + 2 μ l EcoRI
 + 2 μ l Buffer H + H₂O 14.8 μ l, 37°C, 1 hr.

2 μ g/3.2 μ l
 12.8 H₂O

Blunt


+ 10 μ M dNTP (2 μ l (10 μ M/each) + 2 μ l Buffer B

14 μ l H₂O + 4 U/2 μ l Klenow, 25°C, 30 min; + EDTA \rightarrow 10 min

Inactivate Klenow by heating at 75°C, 10 min. Phenol:Chloroform extraction
 for Gene Clean

Then + 2 μ l Not I, 37°C, 1 hr.

③ Run gel, cut bands, pool together,
 Gene Clean. Elute in 40 μ l



pSH1/s-Fv1s1-YM-E
 + Not I + Sac I

④ Ligation

Eluent 16 μ l + 2 μ l 10X Ligation Buffer
 + 2 μ l Taq ligase, 16°C, overnight.


pAdTrack-CMV
 + Not I + EcoRV
 ↓
 pSH1/s-E-
 Fv1-Fv1s1-
 -ICEst
 + Not I
 + Sac I

pSH1/s-
 Fv1s1-
 -YM-E
 + Not I
 + Sac I



⑤ Transformation

⑥ Mini-prep with Qagen kit.

FIG 28



pAdTrack-CMV-SFV/Sl-YM-E



F. G - 29

JPEG image 511x431 pixels







FIG 30

A 3





FIG 31




FIG 32

006260" 8744960

Generation of Ad-GFP-CMV-YM-E
Ad-GFP-CMV-E-ICE

① Co-transform E. coli BJ 5183 with pAdEasy-1 and pAdTrack-CMV-Fv1s1-YM-E or pAdTrack-CMV-E-Fv1-Fv1s1-ICE st.

a. Linearize the shuttle plasmids with Pme I

1 µg of pAdTrack-CMV-Fv1s1-YME

or pAdTrack-CMV-E-Fv1-Fv1s1-ICE st

+ 1 µl Buffer + H₂O → 9 µl, + 1 µl Pme I

37°C, 2 hr.

phenol-chloroform extraction, ethanol precipitation and resuspend in 6 µl H₂O.

b. Co-transformation: with 100 ng pAdEasy-1

20 µl of BJ 5183

2.0 mm cuvettes

at 250V, 200 Ohms, 25 µFD

c. Selection:

pick 10 smallest colonies, miniprep, check with Pac I.

re-transform XLI-Blue with the correct plasmid, miniprep, recheck with Pac I.

Midiprep.

d. Transfect 293 cells with FuGene.

90% confluence 293 cells in 6-well-plate,

4 µg plasmid/6 µl FuGene / well.

FIG 33

U9/647418




FIG 34

006260" 87424960

4 EcoRI + 4



Midiprep
082498




Fig35

pAd Easy-1-Track-CMV-ICE-E
-YAMA-E






FIG 3b



006260" 87424350

FIG 37 Generation of Ad-CMV-E-FV1-FVLS1-ICEst

Construction of pshuttle-CMV-E-FV1-FVLS1-ICEst and Ad-CMV-E-ICE

① cut pshuttle-CMV with EcoRV + NotI

1 μ g / 2 μ l (pshuttle-CMV) + 1 μ l Buffer H
+ 1 μ l EcoRV + 1 μ l NotI + H₂O 5 μ l
37°C, 2hr

② cut pSH/S-E-FV1-FVLS1-ICEst with NotI + EcoRV (get E-FV1-FVLS1-ICEst)

3 μ g / 4 μ l + 3 μ l Buffer H + 3 μ l EcoRV
+ H₂O 21 μ l, 37°C, 1hr

Blunt

+ 10 mM DTT 15 μ l (10 mM / each) + 3 μ l Buffer 2
+ 3 μ l H₂O + 7 μ l T4 DNA Polymerase, 25°C, 30'
+ EDTA \rightarrow 10 mM, 75°C, 10'
phenol:chloroform extraction (twice)
ethanol precipitation \rightarrow 24 μ l H₂O

Then

+ 3 μ l Buffer H + 3 μ l NotI
37°C, 2hr

③ Run Gel, cut bands, pool together, GeneClean II into 4 μ l H₂O

2) Ligation



Elution: 16 μ l + 2 μ l 10x Ligation Buffer
+ 2 μ l T4 Ligase, 25°C overnight


④ Transformation

E: Miniprep + Restriction Enzyme checking

⑦ cut pshuttle-CMV-E-FV1-FVLS1-ICEst with PmeI (1/2 of a miniprep), Cotransform E. coli BJ5183 with
100 ng pAdEasy-1. Miniprep + PacI checking (photo above)

⑧ chose #2, retransform XL-1 Blue, Miniprep, + PacI; Midi prep, + PacI checking (small photo above)









Fig 39

FIG 40

Luciferase Assay of pSiville-CMV

- Fv1 - FvSI - ICE - E

0104 - 010799

- ① 1×10^5 JD-201 cells / well (6 well - plate)
in 1640 RPMI media with 5% FCS
incubate 24 hrs

- ② Transfection: 2 μ g pGL2 2 μ g ICE per well
Fugene 2 μ l / μ g DNA

incubate 6 hrs, add AP1903 at 50 nM.

Incubate 24 hrs.

- ③ lysis - luminometer

8 JAN 99 AM 10:59

V. 1.1

MEAS. TIME(s) : 15.0

SAMPLE RLU

% C.V.

1 1 294

1 2 291

MEAN 292

0.8

Control

2 1 2681959

2 2 2769999

MEAN 2725979

2.3 Control + pGL2

3 1 208762

3 2 223711

MEAN 216237

4.9 pGL2 + ICE

4 1 22084

4 2 22859

MEAN 22471

2.4 pGL2 + ICE + AP1903

006260" 8TH4960

FIG 41

JD-2A, T-C₂, T-C₂G, LNCaP — Adeno-TM or Adeno-ICE

① Plating cells: 24 well-plate

JD-2A — 20000/well

T-C₂ > 10000/wellT-C₂G

LNCaP — 30000/well

in 1ml RPMI 1640 Media (5% FBS)

in 1ml DMEM Media (+5% FBS, 5% Nucserum, Insulin
DHT)

incubate JD 24 hrs

T-C₂, T-C₂G — 48 hrs

LNCaP — 72 hrs

Cell number doubling — 40000

— 40000

— 60000

② Infection:

For JD: ICE 1×10^7 iu/ml

MOI

1

2

4

8

16

Add

4

8

16

32

64 μ lYM 2×10^7 iu/ml

MOI

2.5

5

10

20

40

5

10

20

40

80 μ l

1:10

For T-C₂ICE 5×10^7 iu/ml

MOI

2.5

5

10

20

40

+ T-C₂G

Add

2

4

8

16

32

YM 5×10^7 iu/ml

MOI

2.5

5

10

20

40

Add

2

4

8

16

32

For LNCaP

3

6

12

24

48

After 4 hrs. Add AP 1903 \rightarrow 50 nM, incubate 24 hrs.

③ Fix with 1% Glutaraldehyde — PBS, 15';

Stain 0.5% Crystal violet, 20';

Wash with H₂O, 30';Resolve with Safran's Solution, 200 μ l/well, 5';Transfer 100 μ l to each well in 96-well plate. Read OD at 570 nm.

006260" 2424960

T-C2G Cells Treated with Ad-YAMA or Ad-ICE + AP1903 or AP20187




Fig 42

09/647418




Fig 43




FIG 44




Fig 45

Western Blot 1103-110495

293 cells, infected with Ad-YM or Ad-ICE

9% Resolving Gel; Transfer - 400 mA, 2 hrs.

1:1000 α -HA11 - 1st Ab.

- 1 Ad-YM3
 2 - +1903 (100 nM)
 3 - +2-D-DCB (100 nM)
 4 - +Both

Ad-ICE

- 6 - +1903
 7 - +2-D-DCB
 8 - +Both

Ad-YM4

- 10 - +1903
 11 - +2-D-DCB
 12 - +Both

Expression and Activation of ICE and YM

KODAK SAFETY FILM

KODAK SAFETY FILM

FIG 4b




fig 47

006250" BT4960

ADV-FKBP/ICE effectively kills PC-3 prostate cancer cells

MOI:
AP1903
(50 nM)




Fig 48

ADV-FKBP/ICE effectively kills JD-2a BPH cells

MOI:
AP1903
(50 nM)





Fig 49

Treatment of s.c. prostate adenocarcinoma in situ with CID-inducible caspases



Analysis: resect tumor, H&E, TUNEL, anti-GFP

Fig 50


ADV-FKBP/ICE effectively kills TRAMP-C2 cells in vivo

20X

20X

no ICE
+ CID

ICE
no CID



ICE
no CID

ICE
no CID

10X

40X


Fig 51

*ADV-FKBP/ICE + CID very effectively
kills TRAMP-C2 cells in vivo*

20X

ICE
+ CID

40X

ICE
+ CID

20X

ICE
+ CID

40X


ICE
+ CID

Fig 52

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.